



TRP1 mouse mAb

Catalog No BYmab-18080 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name TYRP1 CAS2 TYRP TYRRP Protein Name (S-6-dilhydroxyindole-2-carboxylic acid oxidase (DHICA oxidase) (EC 1.14.18) (Catalase B) (Glycoprotein 75) (Melanoma antigen gp75) (Tyrosinase-related protein 1) (TRP) (TRP-1) (TRP1) Immunogen Synthesized peptide derived from human TRP1 Specificity This antibody detects endogenous levels of TRP1 at Human, Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane; Single-pass type I membrane protein Melanosome membrane; Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells.<		
Applications WB Gene Name TYRP1 CAS2 TYRP TYRRP Protein Name 5,6-dihydroxyindole-2-carboxylic acid oxidase (DHICA oxidase) (EC 1.14.18) (Catalase B) (Glycoprotein 75) (Melanoma antigen gp75) (Tyrosinase-related protein 1) (TRP) (TRP-1) (TRP) Immunogen Synthesized peptide derived from human TRP1 Specificity This antibody detects endogenous levels of TRP1 at Human, Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane: Single-pass type I membrane protein . Melanosome umembranes: Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB3A, RAB32 and RAB38. Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (i)HICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Ind(2m) Any regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylaring tyrosine and	Catalog No	BYmab-18080
Applications WB Gene Name TYRP1 CAS2 TYRP TYRRP Protein Name \$\begin{array}{cccccccccccccccccccccccccccccccccccc	Isotype	IgG
Gene Name TYRP1 CAS2 TYRP TYRRP Protein Name 5,6-dihydroxyindole-2-carboxylic acid oxidase (DHICA oxidase) (EC 1.14.18) (Catalase B) (Glycoprotein 75) (Melanoma antigen gp75) (Tyrosinase-related protein 1) (TRP1) (TRP-1) (TRP1) Immunogen Synthesized peptide derived from human TRP1 Specificity This antibody detects endogenous levels of TRP1 at Human, Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane : Single-pass type I membrane protein . Melanosome Located to mature stage ill and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Izn(2+) . May regulate or influence the type of melanin synthesized . Also to a lower extent, calMAB1e of hydroxylating tyrosine and	Reactivity	Human;Mouse
Protein Name 5.6-dihydroxyindole-2-carboxylic acid oxidase (DHICA oxidase) (EC 1.14.18-) (Catalase B) (Glycoprotein 75) (Melanoma antigen gp75) (Tyrosinase-related protein 1) (TRP) (TRP-1) (TRP1) Immunogen Synthesized peptide derived from human TRP1 Specificity This antibody detects endogenous levels of TRP1 at Human, Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane : Single-pass type I membrane protein . Melanosome . Located to mature stage Ill and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) . May regulate or influence the type of melanin synthesized . Also to a lower extent, calMABle of hydroxylating tyrosine and	Applications	WB
(Cataláse B)' (Glycoprotein 75) (Melanoma antigen gp75) (Tyrośinase-related protein 1) (TRP) (TRP-1) (TRP-1) Immunogen Synthesized peptide derived from human TRP1 Specificity This antibody detects endogenous levels of TRP1 at Human, Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane : Single-pass type I membrane protein . Melanosome . Located to mature stage III and IV melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dilhydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carb	Gene Name	TYRP1 CAS2 TYRP TYRRP
Specificity This antibody detects endogenous levels of TRP1 at Human, Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane; Single-pass type I membrane protein . Melanosome . Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized . Also to a lower extent, calMABle of hydroxylating tyrosine and	Protein Name	5,6-dihydroxyindole-2-carboxylic acid oxidase (DHICA oxidase) (EC 1.14.18) (Catalase B) (Glycoprotein 75) (Melanoma antigen gp75) (Tyrosinase-related protein 1) (TRP) (TRP-1) (TRP1)
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane: Single-pass type I membrane protein . Melanosome . Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized . Also to a lower extent, caMABle of hydroxylating tyrosine and	Immunogen	Synthesized peptide derived from human TRP1
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane ; Single-pass type I membrane protein . Melanosome . Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) . May regulate or influence the type of melanin synthesized . Also to a lower extent, calMABle of hydroxylating tyrosine and	Specificity	This antibody detects endogenous levels of TRP1 at Human, Mouse
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane : Single-pass type I membrane protein . Melanosome . Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 Tissue Specificity Pigment cells. Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) . May regulate or influence the type of melanin synthesized . Also to a lower extent, calMABle of hydroxylating tyrosine and	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane ; Single-pass type I membrane protein . Melanosome . Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 . Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) . May regulate or influence the type of melanin synthesized . Also to a lower extent, caMABle of hydroxylating tyrosine and	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Melanosome membrane; Single-pass type I membrane protein. Melanosome. Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis. Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 59kD Cell Pathway Melanosome membrane ; Single-pass type I membrane protein . Melanosome . Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 Tissue Specificity Pigment cells. Function Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) . May regulate or influence the type of melanin synthesized . Also to a lower extent, caMABle of hydroxylating tyrosine and	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Observed Band 59kD Melanosome membrane; Single-pass type I membrane protein. Melanosome. Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells. Plays a role in melanin biosynthesis. Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and	Concentration	1 mg/ml
Synonyms Observed Band 59kD Cell Pathway Melanosome membrane; Single-pass type I membrane protein. Melanosome. Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells. Plays a role in melanin biosynthesis. Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and	Purity	≥90%
Observed Band Cell Pathway Melanosome membrane; Single-pass type I membrane protein. Melanosome. Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells. Plays a role in melanin biosynthesis. Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and	Storage Stability	-20°C/1 year
Cell Pathway Melanosome membrane; Single-pass type I membrane protein. Melanosome. Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38. Tissue Specificity Pigment cells. Plays a role in melanin biosynthesis. Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and	Synonyms	
Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 Pigment cells. Plays a role in melanin biosynthesis . Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) . May regulate or influence the type of melanin synthesized . Also to a lower extent, caMABle of hydroxylating tyrosine and	Observed Band	59kD
Function Plays a role in melanin biosynthesis. Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and	Cell Pathway	Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A,
5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and	Tissue Specificity	Pigment cells.
	Function	5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+). May regulate or influence the type of melanin synthesized. Also to a lower extent, caMABle of hydroxylating tyrosine and

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images