



USP22 mouse mAb

Catalog No	BYmab-18198
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	USP22 KIAA1063 USP3L
Protein Name	Ubiquitin carboxyl-terminal hydrolase 22 (EC 3.4.19.12) (Deubiquitinating enzyme 22) (Ubiquitin thioesterase 22) (Ubiquitin-specific-processing protease 22)
Immunogen	Synthesized peptide derived from human USP22
Specificity	This antibody detects endogenous levels of USP22 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year 58kD
Synonyms	·
Synonyms Observed Band	58kD
Synonyms Observed Band Cell Pathway	Nucleus . Moderately expressed in various tissues including heart and skeletal muscle, and weakly expressed in lung and liver. Histone deubiquitinating component of the transcription regulatory histone acetylation (HAT) complex SAGA. Catalyzes the deubiquitination of both histones H2A and H2B, thereby acting as a coactivator. Recruited to specific gene promoters by activators such as MYC, where it is required for transcription.
Synonyms Observed Band Cell Pathway Tissue Specificity	Nucleus . Moderately expressed in various tissues including heart and skeletal muscle, and weakly expressed in lung and liver. Histone deubiquitinating component of the transcription regulatory histone acetylation (HAT) complex SAGA. Catalyzes the deubiquitination of both histones H2A and H2B, thereby acting as a coactivator. Recruited to specific gene promoters by activators such as MYC, where it is required for transcription.
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Nucleus . Moderately expressed in various tissues including heart and skeletal muscle, and weakly expressed in lung and liver. Histone deubiquitinating component of the transcription regulatory histone acetylation (HAT) complex SAGA. Catalyzes the deubiquitination of both histones H2A and H2B, thereby acting as a coactivator. Recruited to specific gene

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd