



NAT1 mouse mAb

Catalog No BYmab-18280 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name NAT1 AAC1 Protein Name Anylamine N-acetyltransferase 1 (EC 2.3.1.5) (Arylamide acetylase 1) (Monomorphic arylamine N-acetyltransferase) (MNAT) (N-acetyltransferase type 1) (NAT-1) Immunogen Synthesized peptide derived from human NAT1 Specificity This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Purity acrease Stability acreas		
Reactivity Human;Mouse;Rat Applications WB Gene Name NAT1 AAC1 Protein Name Arylamine N-acetyltransferase 1 (EC 2.3.1.5) (Arylamide acetylase 1) (Monomorphic arylamine N-acetyltransferase) (MNAT) (N-acetyltransferase type 1) (NAT-1) Immunogen Synthesized peptide derived from human NAT1 Specificity This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bloactivate several known carcinogens.	Catalog No	BYmab-18280
Applications WB Gene Name NAT1 AAC1 Protein Name Avalamine N-acetyltransferase 1 (EC 2.3.1.5) (Arylamide acetylase 1) (Monomorphic arylamine N-acetyltransferase) (MNAT) (N-acetyltransferase type 1) (NAT-1) (Monomorphic arylamine N-acetyltransferase) (MNAT) (N-acetyltransferase type 1) (NAT-1) (MNAT-1) Specificity This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name NAT1 AAC1 Protein Name Arylamine N-acetyltransferase 1 (EC 2.3.1.5) (Arylamide acetylase 1) (Monomorphic arylamine N-acetyltransferase) (MNAT) (N-acetyltransferase type 1) (NAT-1) Immunogen Synthesized peptide derived from human NAT1 Specificity This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Mouse;Rat
Protein Name Arylamine N-acetyltransferase 1 (EC 2.3.1.5) (Arylamide acetylase 1) (Monomorphic arylamine N-acetyltransferase) (MNAT) (N-acetyltransferase type 1) (NAT-1) Immunogen Synthesized peptide derived from human NAT1 Specificity This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Applications	WB
Inmunogen Synthesized peptide derived from human NAT1 Specificity This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Gene Name	NAT1 AAC1
Specificity This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bloactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Protein Name	Arylamine N-acetyltransferase 1 (EC 2.3.1.5) (Arylamide acetylase 1) (Monomorphic arylamine N-acetyltransferase) (MNAT) (N-acetyltransferase type 1) (NAT-1)
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bloactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human NAT1
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of NAT1 at Human, Mouse,Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background Matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background Matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 32kD Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background Matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm. Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background Matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background Avoid repeated freezing and thawing!	Observed Band	32kD
Function Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background Avoid repeated freezing and thawing!	Cell Pathway	Cytoplasm.
Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens. Background Matters needing Avoid repeated freezing and thawing!	Tissue Specificity	
matters needing Avoid repeated freezing and thawing!	Function	Participates in the detoxification of a plethora of hydrazine and arylamine drugs. Catalyzes the N- or O-acetylation of various arylamine and heterocyclic amine substrates and is able to bioactivate several known carcinogens.
matters needing Avoid repeated freezing and thawing! attention	Background	
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd