



APOA rabbit pAb

Catalog No	BYab-00781
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB; ELISA
Gene Name	LPA
Protein Name	APOA
Immunogen	Synthesized peptide derived from human APOA
Specificity	This antibody detects endogenous levels of Human APOA
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Apolipoprotein(a);Apo(a);Lp(a);EC 3.4.21.-)
Observed Band	
Cell Pathway	
Tissue Specificity	
Function	disease:Elevated plasma concentrations of apo(a) and its naturally occurring proteolytic fragments are correlated with atherosclerosis. Homology with plasminogen kringle IV and V is thought to underlie the atherogenicity of the protein, because the fragments are competing with plasminogen for fibrin(ogen) binding.,function:Apo(a) is the main constituent of lipoprotein(a) (Lp(a)). It has serine proteinase activity and is able of autoproteolysis. Inhibits tissue-type plasminogen activator 1. Lp(a) may be a ligand for megalin/Gp 330.,miscellaneous:Apo(a) is known to be proteolytically cleaved, leading to the formation of the so-called mini-Lp(a). Apo(a) fragments accumulate in atherosclerotic lesions, where they may promote thrombogenesis. O-glycosylation may limit the extent of proteolytic fragmentation.,online information:The Singapore human mutation and polymorphism

Nanjing BYabscience technology Co.,Ltd



database,polymorphis

Background

The protein encoded by this gene is a serine proteinase that inhibits the activity of tissue-type plasminogen activator I. The encoded protein constitutes a substantial portion of lipoprotein(a) and is proteolytically cleaved, resulting in fragments that attach to atherosclerotic lesions and promote thrombogenesis. Elevated plasma levels of this protein are linked to atherosclerosis. Depending on the individual, the encoded protein contains 2-43 copies of kringle-type domains. The allele represented here contains 15 copies of the kringle-type repeats and corresponds to that found in the reference genome sequence. [provided by RefSeq, Dec 2009],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images**Nanjing BYabscience technology Co.,Ltd**网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658