



Ikaros Monoclonal Antibody

Catalog No	BYab-01043
Isotype	IgG
Reactivity	Human;Mouse;Dog
Applications	WB
Gene Name	IKZF1
Protein Name	DNA-binding protein Ikaros
Immunogen	Purified recombinant human Ikaros (C-terminus) protein fragments expressed in E.coli.
Specificity	Ikaros Monoclonal Antibody detects endogenous levels of Ikaros protein.
Formulation	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/1000 - 1/2000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	IKZF1; IK1; IKAROS; LYF1; ZNFN1A1; DNA-binding protein Ikaros; Ikaros family zinc finger protein 1; Lymphoid transcription factor LyF-1
Observed Band	
Cell Pathway	Nucleus . In resting lymphocytes, distributed diffusely throughout the nucleus. Localizes to pericentromeric heterochromatin in proliferating cells. This localization requires DNA binding which is regulated by phosphorylation / dephosphorylation events. .; [Isoform Ik2]: Nucleus. In resting lymphocytes, distributed diffusely throughout the nucleus. Localizes to pericentromeric heterochromatin in proliferating cells. This localization requires DNA binding which is regulated by phosphorylation / dephosphorylation events (By similarity). .; [Isoform Ik6]: Cytoplasm .
Tissue Specificity	Abundantly expressed in thymus, spleen and peripheral blood Leukocytes and lymph nodes. Lower expression in bone marrow and small intestine.
Function	function: Binds and activates the enhancer (delta-A element) of the CD3-delta gene. Functions in the specification and the maturation of the T-lymphocyte. Also interacts with a critical control element in the TDT (terminal deoxynucleotidyltransferase) promoter as well as with the promoters for other

Nanjing BYabs science technology Co.,Ltd



genes expressed during early stages of B- and T-cell development.,similarity:Belongs to the Ikaros C2H2-type zinc-finger protein family.,similarity:Contains 6 C2H2-type zinc fingers.,subunit:Interacts with IKZF4 AND IKZF5.,tissue specificity:Abundantly expressed in thymus, spleen and peripheral blood Leukocytes and lymph nodes. Lower expression in bone marrow and small intestine.,

Background

This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using Ikaros Monoclonal Antibody against Ramos, Jurkat cell lysate.