



MyoD (phospho Ser200) Polyclonal Antibody

Catalog No	BYab-01296
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	MYOD1
Protein Name	Myoblast determination protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from human MYOD around the phosphorylation site of Ser200. AA range:171-220
Specificity	Phospho-MyoD (S200) Polyclonal Antibody detects endogenous levels of MyoD protein only when phosphorylated at S200.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MYOD1; BHLHC1; MYF3; MYOD; Myoblast determination protein 1; Class C basic helix-loop-helix protein 1; bHLHc1; Myogenic factor 3; Myf-3
Observed Band	34kD
Cell Pathway	Nucleus.
Tissue Specificity	Muscle,Skeletal muscle,
Function	function:Involved in muscle differentiation (myogenic factor). Induces fibroblasts to differentiate into myoblasts. Activates muscle-specific promoters. Interacts with and is inhibited by the twist protein. This interaction probably involves the basic domains of both proteins.,online information:MyoD entry,PTM:Acetylated by a complex containing EP300 and PCAF. The acetylation is essential to activate target genes. Conversely, its deacetylation by SIRT1 inhibits its function.,PTM:Ubiquitinated on the N-terminus; which is required for proteasomal degradation.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Seems to form active heterodimers with ITF-2. Interacts with SUV39H1.,


Nanjing BYabscience technology Co.,Ltd




Background	This gene encodes a nuclear protein that belongs to the basic helix-loop-helix family of transcription factors and the myogenic factors subfamily. It regulates muscle cell differentiation by inducing cell cycle arrest, a prerequisite for myogenic initiation. The protein is also involved in muscle regeneration. It activates its own transcription which may stabilize commitment to myogenesis. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Enzyme-Linked Immunosorbent Assay
(Phospho-ELISA) for Immunogen Phosphopeptide
(Phospho-left) and Non-Phosphopeptide
(Phospho-right), using MYOD (Phospho-Ser200)
Antibody



Western blot analysis of lysates from Jurkat cells
treated with Ca⁺ 40nM 30', using MYOD
(Phospho-Ser200) Antibody. The lane on the right is
blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded
human spleen. 1, Tris-EDTA, pH9.0 was used for
antigen retrieval. 2 Antibody was diluted at 1:200(4°
overnight). 3, Secondary antibody was diluted at
1:200(room temperature, 45min).

Nanjing BYabs science technology Co.,Ltd