



ATRX Polyclonal Antibody

Catalog No	BYab-01558
Isotype	IgG
Reactivity	Human;Mouse
Applications	IF;ELISA
Gene Name	ATRX
Protein Name	Transcriptional regulator ATRX
Immunogen	The antiserum was produced against synthesized peptide derived from human ATRX. AA range:111-160
Specificity	ATRX Polyclonal Antibody detects endogenous levels of ATRX protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ATRX; RAD54L; XH2; Transcriptional regulator ATRX; ATP-dependent helicase ATRX; X-linked helicase II; X-linked nuclear protein; XNP; Znf-HX
Observed Band	
Cell Pathway	Nucleus. Chromosome, telomere. Nucleus, PML body. Associated with pericentromeric heterochromatin during interphase and mitosis, probably by interacting with CBX5/HP1 alpha. Colocalizes with histone H3.3, DAXX, HIRA and ASF1A at PML-nuclear bodies. Colocalizes with cohesin (SMC1 and SMC3) and MECP2 at the maternal H19 ICR (By similarity).
Tissue Specificity	Ubiquitous.
Function	disease:Defects in ATRX are a cause of alpha-thalassemia myelodysplasia syndrome (ATMDS) [MIM:300448]. In this disorder, alpha-thalassemia occurs as an acquired abnormality in association with a multilineage myelodysplasia.,disease:Defects in ATRX are the cause of mental retardation syndromic X-linked with hypotonic facies syndrome type 1 (MRXSHF1) [MIM:309580]; also called Carpenter-Waziri syndrome (CWS), Juberg-Marsidi syndrome (JMS), Smith-Fineman-Myers syndrome type 1 (SFM1). Clinical

Nanjing BYabs science technology Co.,Ltd



features include severe mental retardation, dysmorphic facies, and a highly skewed X-inactivation pattern in carrier women. Other more variable features include hypogonadism, deafness, renal anomalies, and mild skeletal defects. disease: Defects in ATRX are the cause of X-linked alpha-thalassemia/mental retardation syndrome (ATR-X) [MIM:301040]. ATR-X is an X-linked disorder comprising severe psychomotor

Background

ATRX, chromatin remodeler(ATRX) Homo sapiens The protein encoded by this gene contains an ATPase/helicase domain, and thus it belongs to the SWI/SNF family of chromatin remodeling proteins. This protein is found to undergo cell cycle-dependent phosphorylation, which regulates its nuclear matrix and chromatin association, and suggests its involvement in the gene regulation at interphase and chromosomal segregation in mitosis. Mutations in this gene are associated with an X-linked mental retardation (XLMR) syndrome most often accompanied by alpha-thalassemia (ATRX) syndrome. These mutations have been shown to cause diverse changes in the pattern of DNA methylation, which may provide a link between chromatin remodeling, DNA methylation, and gene expression in developmental processes. Multiple alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Aug 2013],



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of A549 cells, using ATRX Antibody. The picture on the right is blocked with the synthesized peptide.