



DDX8 Polyclonal Antibody

Catalog No	BYab-01648
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	DHX8
Protein Name	ATP-dependent RNA helicase DHX8
Immunogen	The antiserum was produced against synthesized peptide derived from human DHX8. AA range:391-440
Specificity	DDX8 Polyclonal Antibody detects endogenous levels of DDX8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DHX8; DDX8; ATP-dependent RNA helicase DHX8; DEAH box protein 8; RNA helicase HRH1
Observed Band	150kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in skin, cervix and nerve. Also expressed in the brain.
Function	domain:The RS domain confers a nuclear localization signal, and appears to facilitate the interaction with the spliceosome.,function:Facilitates nuclear export of spliced mRNA by releasing the RNA from the spliceosome.,similarity:Belongs to the DEAD box helicase family. DEAH subfamily. DDX8/PRP22 sub-subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 S1 motif domain.,subunit:Identified in the spliceosome C complex, at least composed of AQR, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38, DHX8, EFTUD2, FRG1, GPATC1, HNRPA1, HNRPA2B1, HNRPA3, HNRPC, HNRPF, HNRPH1, HNRPK, HNRPM, HNRPR,


Nanjing BYabscience technology Co.,Ltd




	HNRPU, KIAA1160, KIAA1604, LSM2, LSM3, MAGOH, MORG1, PABPC1, PLRG1, PNN, PPIE, PPIL1, PPIL3, PPWD1, PRPF19, PRPF4B, PRPF6, PRPF8, RALY, RBM22, RBM8A, RBMX, SART1, SF3A1, SF3A2, SF3A3, SF3B1, SF3
Background	This gene is a member of the DEAH box polypeptide family. The encoded protein contains the DEAH (Asp-Glu-Ala-His) motif which is characteristic of all DEAH box proteins, and is thought to function as an ATP-dependent RNA helicase that regulates the release of spliced mRNAs from spliceosomes prior to their export from the nucleus. This protein may be required for the replication of human immunodeficiency virus type 1 (HIV-1). Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using DDX8 Polyclonal Antibody diluted at 1:500 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Western blot analysis of the lysates from HeLa cells using DHX8 antibody.