



Nanog P8 Polyclonal Antibody

Catalog No	BYab-01890
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IF;ELISA;IHC
Gene Name	NANOGP8
Protein Name	Putative homeobox protein NANOGP8
Immunogen	The antiserum was produced against synthesized peptide derived from human NANOGP8. AA range:51-100
Specificity	Nanog P8 Polyclonal Antibody detects endogenous levels of Nanog P8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000; IF/ICC 1:50-200;ELISA 1:2000-20000;IHC-p 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NANOGP8; Putative homeobox protein NANOGP8
Observed Band	35kD
Cell Pathway	Nucleus .
Tissue Specificity	Embryonic stem cell,Teratocarcinoma,Urinary bladder carcinoma,
Function	developmental stage:Expressed in embryonic stem (ES) and carcinoma (EC) cells. Expressed in inner cell mass (ICM) of the blastocyst and gonocytes between 14 and 19 weeks of gestation (at protein level). Not expressed in oocytes, unfertilized oocytes, 2-16 cell embryos and early morula (at protein level). Expressed in embryonic stem cells (ES). Expression decreases with ES differentiation.,function:May act as a transcription regulator (By similarity). When overexpressed, promotes cells to enter into S phase and proliferation.,function:Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophoctoderm lineages. Blocks bone morphogenetic protein-induced mesoderm

Nanjing BYabscience technology Co.,Ltd



differentiation of ES cells by physically interacting with SMAD1 an

Background

This locus is a processed pseudogene of the transcription factor NANOG. NANOG plays a central role in regulating self-renewal in pluripotent stem cells and tumor cells. This pseudogene contains an intact open reading frame that could potentially encode a protein similar to NANOG. Although there is no evidence of transcription from this pseudogene, RT-PCR studies suggest that NANOGP8 may be expressed in some cancer cell lines. In vitro studies using a recombinant NANOGP8 protein have shown that the protein localizes to the nucleus and can promote cell proliferation, similar to NANOG. [provided by RefSeq, Sep 2009],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd




Products Images



Western Blot analysis of various cells using Nanog P8 Polyclonal Antibody diluted at 1:2000



Western blot analysis of lysates from HUVEC, HT-29, HepG2, and Jurkat cells, using NANOGP8 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658