



SMUG1 Polyclonal Antibody

Catalog No	BYab-02023
Isotype	IgG
Reactivity	Human;Mouse;Rat;Monkey
Applications	WB;IHC;IF;ELISA
Gene Name	SMUG1
Protein Name	Single-strand selective monofunctional uracil DNA glycosylase
Immunogen	The antiserum was produced against synthesized peptide derived from human SMUG1. AA range:91-140
Specificity	SMUG1 Polyclonal Antibody detects endogenous levels of SMUG1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SMUG1; Single-strand selective monofunctional uracil DNA glycosylase
Observed Band	32kD
Cell Pathway	Nucleus .
Tissue Specificity	Adipose tissue,Brain,Liver,Muscle,Ovary,Ute
Function	function:Responsible for recognizing base lesions in the genome and initiating base excision DNA repair. Acts as a monofunctional DNA glycosylase specific for uracil (U) residues in DNA and has a preference for single-stranded DNA substrates. The activity is greater against mismatches (U/G) than against matches (U/A). Excised uracil (U), 5-formyluracil (fU) and uracil derivatives bearing an oxidized group at C5 [5-hydroxyuracil (hoU) and 5-hydroxymethyluracil (hmU)] in ssDNA and dsDNA but not analogous cytosine derivatives (5-hydroxycytosine and 5-formylcytosine) and other oxidized damage. The activity is damage specificity and salt concentration-dependent. The general order of the preference for ssDNA and dsDNA is the following: ssDNA > dsDNA (G pair) = dsDNA (A pair) at the low salt concentration. At the high concentration dsDNA (G pair) > dsDNA

Nanjing BYabscience technology Co.,Ltd



(A pair) > ssDNA.,

Background

This gene encodes a protein that participates in base excision repair by removing uracil from single- and double-stranded DNA. Many alternatively spliced transcript variants exist for this gene; the full-length nature is known for some but not all of the variants. [provided by RefSeq, Aug 2011],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658