



YAP Polyclonal Antibody

Catalog No	BYab-02158
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	YAP1
Protein Name	Yorkie homolog
Immunogen	The antiserum was produced against synthesized peptide derived from human YAP. AA range:281-330
Specificity	YAP Polyclonal Antibody detects endogenous levels of YAP protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	YAP1; YAP65; Yorkie homolog; 65 kDa Yes-associated protein; YAP65
Observed Band	67kD
Cell Pathway	Cytoplasm . Nucleus . Both phosphorylation and cell density can regulate its subcellular localization (PubMed:18158288, PubMed:20048001). Phosphorylation sequesters it in the cytoplasm by inhibiting its translocation into the nucleus (PubMed:18158288, PubMed:20048001). At low density, predominantly nuclear and is translocated to the cytoplasm at high density (PubMed:18158288, PubMed:20048001, PubMed:25849865). PTPN14 induces translocation from the nucleus to the cytoplasm (PubMed:22525271). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity). .
Tissue Specificity	Increased expression seen in some liver and prostate cancers. Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease (at protein level).
Function	PTM:Phosphorylated upon DNA damage, probably by ATM or


Nanjing BYabscience technology Co.,Ltd




	ATR.,similarity:Contains 1 WW domain.,subunit:Binds to the SH3 domain of the YES kinase. Binds to WBP1 and WBP2. Binds, in vitro, through the WW1 domain, to neural isoforms of ENAH that contain the PPSY motif.,
Background	This gene encodes a downstream nuclear effector of the Hippo signaling pathway which is involved in development, growth, repair, and homeostasis. This gene is known to play a role in the development and progression of multiple cancers as a transcriptional regulator of this signaling pathway and may function as a potential target for cancer treatment. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Aug 2013],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western blot analysis of lysates from 1)Hela cell ,
2)Hela cells knockdown by siRNA
(F:GGUCAGAGAUACUUCUUAATT,R:UUAAGAAGU
AUCUCUGACCTT), (Green) primary antibody was
diluted at 1:1000, 4° over night, Dylight 800 secondary
antibody(Immunoway:RS23920)was diluted at
1:10000, 37° 1hour. (Red) GAPDH Monoclonal
Antibody(5B7) (Immunoway:YM3029) antibody was
diluted at 1:5000 as loading control, 4° over night,
Dylight 680 secondary
antibody(Immunoway:RS23710)was diluted at
1:10000, 37° 1hour.



Immunofluorescence analysis of Hela cell. 1,YAP
Polyclonal Antibody(red) was diluted at 1:200(4°
overnight). GAPDH Monoclonal Antibody(2B8)(green)
was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit
Alexa Fluor 594 Catalog:RS3611 was diluted at
1:1000(room temperature, 50min). Goat Anti Mouse
Alexa Fluor 488 Catalog:RS3208 was diluted at
1:1000(room temperature, 50min).




A

B

C

Immunofluorescence analysis of rat-spleen tissue.
1,YAP Polyclonal Antibody(red) was diluted at 1:200(4°
C,overnight). 2, Cy3 labeled Secondary antibody was
diluted at 1:300(room temperature, 50min).3, Picture B:
DAPI(blue) 10min. Picture A:Target. Picture B: DAPI.
Picture C: merge of A+B




A

B

C

Immunofluorescence analysis of rat-spleen tissue.
1,YAP Polyclonal Antibody(red) was diluted at 1:200(4°
C,overnight). 2, Cy3 labeled Secondary antibody was
diluted at 1:300(room temperature, 50min).3, Picture B:
DAPI(blue) 10min. Picture A:Target. Picture B: DAPI.
Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Human-uterus tissue. 1,YAP Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.