



MAN1B1 Polyclonal Antibody

Catalog No	BYab-02666
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	MAN1B1
Protein Name	Endoplasmic reticulum mannosyl-oligosaccharide 1,2-alpha-mannosidase
Immunogen	Synthesized peptide derived from MAN1B1 . at AA range: 100-180
Specificity	MAN1B1 Polyclonal Antibody detects endogenous levels of MAN1B1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MAN1B1; Endoplasmic reticulum mannosyl-oligosaccharide 1; 2-alpha-mannosidase; ER alpha-1,2-mannosidase; ER mannosidase 1; ERMan1; Man9GlcNAc2-specific-processing alpha-mannosidase; Mannosidase alpha class 1B member 1
Observed Band	80kD
Cell Pathway	Endoplasmic reticulum membrane ; Single-pass type II membrane protein .
Tissue Specificity	Widely expressed.
Function	catalytic activity:Hydrolysis of the terminal (1->2)-linked alpha-D-mannose residues in the oligo-mannose oligosaccharide Man(9)(GlcNAc)(2).,cofactor:Calcium.,enzyme regulation:Inhibited by both 1-deoxymannojirimycin and kifunensine.,function:Involved in the maturation of Asn-linked oligosaccharides. Trim a single alpha-1,2-linked mannose residue from Man(9)GlcNAc(2) to produce Man(8)GlcNAc(2). The only product is the Man(8)GlcNAc(2) isomer B, the form lacking the middle-arm terminal alpha 1,2-mannose. It may be involved in glycoprotein quality control since it is


Nanjing BYabscience technology Co.,Ltd




	important to target misfolded glycoproteins for degradation.,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyl hydrolase 47 family.,tissue specificity:Widely expressed.,
Background	This gene encodes an enzyme belonging to the glycosyl hydrolase 47 family. This enzyme functions in N-glycan biosynthesis, and is a class I alpha-1,2-mannosidase that specifically converts Man9GlcNAc to Man8GlcNAc isomer B. It is required for N-glycan trimming to Man5-6GlcNAc2 in the endoplasmic-reticulum-associated degradation pathway. Mutations in this gene cause autosomal-recessive intellectual disability. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 11. [provided by RefSeq, Dec 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of 3T3 cells using MAN1B1 Polyclonal Antibody



Western blot analysis of MAN1B1 Antibody. The lane on the right is blocked with the MAN1B1 peptide.