



PAH Polyclonal Antibody

Catalog No	BYab-02732
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	PAH
Protein Name	Phenylalanine-4-hydroxylase
Immunogen	The antiserum was produced against synthesized peptide derived from human PAH. AA range:351-400
Specificity	PAH Polyclonal Antibody detects endogenous levels of PAH protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PAH; Phenylalanine-4-hydroxylase; PAH; Phe-4-monooxygenase
Observed Band	51kD
Cell Pathway	cytosol,extracellular exosome,
Tissue Specificity	Liver,
Function	catalytic activity:L-phenylalanine + tetrahydrobiopterin + O(2) = L-tyrosine + 4a-hydroxytetrahydrobiopterin.,cofactor:Fe(2+) ion.,disease:Defects in PAH are the cause of hyperphenylalaninemia (HPA) [MIM:261600]. HPA is the mildest form of phenylalanine hydroxylase deficiency.,disease:Defects in PAH are the cause of non-phenylketonuria hyperphenylalaninemia (Non-PKU HPA) [MIM:261600]. Non-PKU HPA is a mild form of phenylalanine hydroxylase deficiency characterized by phenylalanine levels persistently below 600 mumol, which allows normal intellectual and behavioral development without treatment. Non-PKU HPA is usually caused by the combined effect of a mild hyperphenylalaninemia mutation and a severe one.,disease:Defects in PAH are the cause of phenylketonuria (PKU) [MIM:261600]. PKU is an autosomal

Nanjing BYabscience technology Co.,Ltd



recessive inborn error of phenylalanine metabolism, due to severe phenylalanine hydroxylase

Background

PAH encodes the enzyme phenylalanine hydroxylase that is the rate-limiting step in phenylalanine catabolism. Deficiency of this enzyme activity results in the autosomal recessive disorder phenylketonuria. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PAH Polyclonal Antibody




Immunohistochemistry analysis of PAH antibody in paraffin-embedded human brain tissue.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



Western blot analysis of lysate from HepG2 cells, using PAH antibody.