



Troponin I-C Monoclonal Antibody

Catalog No	BYab-02943
Isotype	IgG
Reactivity	Human
Applications	WB;IHC;IF;ELISA
Gene Name	TNNI3
Protein Name	Troponin I cardiac muscle
Immunogen	Purified recombinant fragment of Troponin I-C expressed in E. Coli.
Specificity	Troponin I-C Monoclonal Antibody detects endogenous levels of Troponin I-C protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	WB: 1/500 - 1/2000. IHC: 1/200 - 1/1000. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TNNI3; TNNC1; Troponin I; cardiac muscle; Cardiac troponin I
Observed Band	
Cell Pathway	cytosol,troponin complex,sarcomere,
Tissue Specificity	Heart,Heart muscle,PCR rescued clones,
Function	disease:Defects in TNNI3 are the cause of cardiomyopathy dilated type 2A (CMD2A) [MIM:611880]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death.,disease:Defects in TNNI3 are the cause of cardiomyopathy familial hypertrophic type 7 (CMH7) [MIM:191044]. Familial hypertrophic cardiomyopathy is a hereditary heart disorder characterized by ventricular hypertrophy, which is usually asymmetric and often involves the interventricular septum. The symptoms include dyspnea, syncope, collapse, palpitations, and chest pain. They can be readily provoked by exercise. The disorder has inter- and intrafamilial variability ranging from benign to malignant forms with high risk of cardiac failure and sudden cardiac death.,disease:Defects in TNNI3 are the cau

Nanjing BYabscience technology Co.,Ltd



Background

Troponin I (TnI), along with troponin T (TnT) and troponin C (TnC), is one of 3 subunits that form the troponin complex of the thin filaments of striated muscle. TnI is the inhibitory subunit; blocking actin-myosin interactions and thereby mediating striated muscle relaxation. The TnI subfamily contains three genes: TnI-skeletal-fast-twitch, TnI-skeletal-slow-twitch, and TnI-cardiac. This gene encodes the TnI-cardiac protein and is exclusively expressed in cardiac muscle tissues. Mutations in this gene cause familial hypertrophic cardiomyopathy type 7 (CMH7) and familial restrictive cardiomyopathy (RCM). [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using Troponin I-C Monoclonal Antibody against truncated Troponin I-C recombinant protein.



Immunohistochemistry analysis of paraffin-embedded human normal cardiac muscle tissue, showing cytoplasmic localization with DAB staining using Troponin I-C Monoclonal Antibody.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658