



SF-1 Polyclonal Antibody

Catalog No	BYab-03336
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	NR5A1
Protein Name	Steroidogenic factor 1
Immunogen	The antiserum was produced against synthesized peptide derived from human NR5A1. AA range:169-218
Specificity	SF-1 Polyclonal Antibody detects endogenous levels of SF-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NR5A1; AD4BP; FTZF1; SF1; Steroidogenic factor 1; SF-1; STF-1; Adrenal 4-binding protein; Fushi tarazu factor homolog 1; Nuclear receptor subfamily 5 group A member 1; Steroid hormone receptor Ad4BP
Observed Band	50kD
Cell Pathway	Nucleus .
Tissue Specificity	High expressed in the adrenal cortex, the ovary, the testis, and the spleen (PubMed:9177385).
Function	disease:Defects in NR5A1 are a cause of adrenocortical insufficiency without ovarian defect [MIM:184757]. The disease is characterized by severe 'slackness,' muscular hypotonia. There is decreased sodium, increased potassium and elevated ACTH.,disease:Defects in NR5A1 are a cause of XY sex reversal with or without adrenal failure [MIM:184757]. This disease is characterized by normal female external genitalia and retention of the uterus.,function:Transcriptional activator. Seems to be essential for sexual differentiation and formation of the primary steroidogenic tissues. Binds to the Ad4 site found in the promoter region of steroidogenic P-450 genes such as CYP11A, CYP11B and CYP21B. Also

Nanjing BYabscience technology Co.,Ltd



regulates the Muellerian inhibiting substance (AMH) gene as well as the AHCH and STAR genes. 5'-YCAAGGYC-3' and 5'-RRAGGTCA-3' are the consensus sequences for the recognition by NR5A1/FTZF1. The SFPQ-NO

Background

The protein encoded by this gene is a transcriptional activator involved in sex determination. The encoded protein binds DNA as a monomer. Defects in this gene are a cause of XY sex reversal with or without adrenal failure as well as adrenocortical insufficiency without ovarian defect. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from rat lung, using STF-1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658