

Peroxin 1 Polyclonal Antibody

Catalog No	BYab-04065
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF;ELISA
Gene Name	PEX1
Protein Name	Peroxisome biogenesis factor 1
Immunogen	The antiserum was produced against synthesized peptide derived from human PEX1. AA range:1234-1283
Specificity	Peroxin 1 Polyclonal Antibody detects endogenous levels of Peroxin 1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PEX1; Peroxisome biogenesis factor 1; Peroxin-1; Peroxisome biogenesis disorder protein 1
Observed Band	
Cell Pathway	Cytoplasm. Peroxisome membrane. Associated with peroxisomal membranes.
Tissue Specificity	Brain,Lymph,Trachea,
Function	disease:Defects in PEX1 are a cause of adrenoleukodystrophy neonatal (NALD) [MIM:202370]. NALD is a peroxisome biogenesis disorder (PBD) characterized by the accumulation of very long-chain fatty acids, adrenal insufficiency and mental retardation.,disease:Defects in PEX1 are a cause of infantile Refsum disease (IRD) [MIM:266510]. IRD is a mild peroxisome biogenesis disorder (PBD). Clinical features include early onset, mental retardation, minor facial dysmorphism, retinopathy, sensorineural hearing deficit, hepatomegaly, osteoporosis, failure to thrive, and hypocholesterolemia. The biochemical abnormalities include accumulation of phytanic acid, very long chain fatty acids (VLCFA), di- and trihydroxycholestanoic acid and pipecolic acid.,disease:Defects in PEX1 are the

Nanjing BYabs science technology Co.,Ltd



cause of peroxisome biogenesis disorder complementation group 1 (PBD-CG1) [MIM:602136]; also known as PBD-CGE. PBD refer

Background

This gene encodes a member of the AAA ATPase family, a large group of ATPases associated with diverse cellular activities. This protein is cytoplasmic but is often anchored to a peroxisomal membrane where it forms a heteromeric complex and plays a role in the import of proteins into peroxisomes and peroxisome biogenesis. Mutations in this gene have been associated with complementation group 1 peroxisomal disorders such as neonatal adrenoleukodystrophy, infantile Refsum disease, and Zellweger syndrome. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PEX1 Antibody. The picture on the right is blocked with the synthesized peptide.