



PPAR Delta mouse Monoclonal Antibody(2F9)

Catalog No	BYab-04842
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	IF;IHC
Gene Name	PPARD NR1C2 PPARB
Protein Name	PPAR Delta
Immunogen	Recombinant Protein of PPAR Delta of PPAR Delta
Specificity	PPAR Delta protein detects endogenous levels of PPAR Delta
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IF: 1:50-200 IHC 1:100-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Peroxisome proliferator-activated receptor delta (PPAR-delta) (NUC1) (Nuclear hormone receptor 1) (NUC1) (Nuclear receptor subfamily 1 group C member 2) (Peroxisome proliferator-activated receptor beta) (PPAR-beta)
Observed Band	50kD
Cell Pathway	Nucleus.
Tissue Specificity	Ubiquitous with maximal levels in placenta and skeletal muscle.
Function	function:Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Decreases expression of NPC1L1 once activated by a ligand.,online information:Peroxisome proliferator-activated receptor entry,similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Heterodimer with the retinoid X receptor.,tissue specificity:Ubiquitous with maximal levels in placenta and skeletal muscle.,

Nanjing BYabscience technology Co.,Ltd

**Background**

peroxisome proliferator activated receptor delta(PPARD) Homo sapiens This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) family. PPARs are nuclear hormone receptors that bind peroxisome proliferators and control the size and number of peroxisomes produced by cells. PPARs mediate a variety of biological processes, and may be involved in the development of several chronic diseases, including diabetes, obesity, atherosclerosis, and cancer. This protein is a potent inhibitor of ligand-induced transcription activity of PPAR alpha and PPAR gamma. It may function as an integrator of transcription repression and nuclear receptor signaling. The expression of this gene is found to be elevated in colorectal cancer cells. The elevated expression can be repressed by adenomatosis polyposis coli (APC), a tumor suppressor protein related to APC/beta-catenin signaling pathway. Knockout studies in mice suggested the role of this

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images




A

B

C

Immunofluorescence analysis of rat-spleen tissue. 1,PPAR Delta Mouse Monoclonal Antibody(2F9)(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




A

B


C

Immunofluorescence analysis of mouse-spleen tissue. 1,PPAR Delta Mouse Monoclonal Antibody(2F9)(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Negative Control

Immunohistochemical analysis of paraffin-embedded Human-uterus tissue. 1,PPAR Delta Mouse Monoclonal Antibody(2F9) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Human Brain Tissue using PPAR Delta Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Mouse Brain Tissue using PPAR Delta Mouse mAb diluted at 1:200.