



CDKL5 Polyclonal Antibody

Catalog No	BYab-04961
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	CDKL5 STK9
Protein Name	Cyclin-dependent kinase-like 5 (EC 2.7.11.22) (Serine/threonine-protein kinase 9)
Immunogen	Synthesized peptide derived from human protein . at AA range: 30-110
Specificity	CDKL5 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	113kD
Cell Pathway	Nucleus . Cytoplasm, cytoskeleton, cilium basal body . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .
Tissue Specificity	Expressed in brain, lung, kidney, prostate, ovary, placenta, pancreas and testis.; [Isoform 2]: Predominant transcript in brain.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is uncertain whether Met-1 or Met-10 is the initiator.,disease:Chromosomal aberrations involving CDKL5 are a cause of X-linked infantile spasm syndrome (ISSX) [MIM:308350]; also known as X-linked West syndrome. Translocation t(X;6)(p22.3;q14); translocation t(X;7)(p22.3;p15). ISSX is characterized by infantile spasms, hypsarrhythmia on EEG, and developmental arrest leading to severe to profound mental retardation.,disease:Defects in CDKL5 are a cause of atypical CDKL5-related Rett syndrome [MIM:300672]. Rett syndrome is an X-linked dominant disease. It is a progressive neurologic developmental disorder and one of the most common causes of mental retardation in females. Patients appear to develop normally until 6 to 18 months of age, then gradually lose

Nanjing BYabscience technology Co.,Ltd



speech and purposeful hand movements and develop microcephaly, se

Background

This gene is a member of Ser/Thr protein kinase family and encodes a phosphorylated protein with protein kinase activity. Mutations in this gene have been associated with X-linked infantile spasm syndrome (ISSX), also known as X-linked West syndrome, and Rett syndrome (RTT). Alternate transcriptional splice variants have been characterized. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658