



CRLF1 Polyclonal Antibody

Catalog No	BYab-05061
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	CRLF1 UNQ288/PRO327
Protein Name	Cytokine receptor-like factor 1 (Cytokine-like factor 1) (CLF-1) (ZcytoR5)
Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Specificity	CRLF1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	46kD
Cell Pathway	Secreted .
Tissue Specificity	Highest levels of expression observed in spleen, thymus, lymph node, appendix, bone marrow, stomach, placenta, heart, thyroid and ovary. Strongly expressed also in fetal lung.
Function	disease:Defects in CRLF1 are the cause of cold-induced sweating syndrome 1 (CISS1) [MIM:272430]. Cold-induced sweating syndrome (CISS) is an autosomal recessive disorder characterized by profuse sweating induced by cool surroundings (temperatures of 7 to 18 degrees Celsius). Additional abnormalities include a high-arched palate, nasal voice, depressed nasal bridge, inability to fully extend the elbows and kyphoscoliosis.,disease:Defects in CRLF1 are the cause of Crisponi syndrome [MIM:601378]. Crisponi syndrome is a rare autosomal recessive disorder characterized by congenital muscular contractions of facial muscles, with trismus in response to stimuli, dysmorphic features, bilateral camptodactyly, major feeding and respiratory difficulties, and access of hyperthermia leading to death in the first months of life.,domain:The WSXWS

Nanjing BYabscience technology Co.,Ltd



motif appears to be necessary for proper protein folding a

Background

This gene encodes a member of the cytokine type I receptor family. The protein forms a secreted complex with cardiotrophin-like cytokine factor 1 and acts on cells expressing ciliary neurotrophic factor receptors. The complex can promote survival of neuronal cells. Mutations in this gene result in Crisponi syndrome and cold-induced sweating syndrome. [provided by RefSeq, Oct 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658