



KDM5A Polyclonal Antibody

Catalog No	BYab-05113
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	KDM5A JARID1A RBBP2 RBP2
Protein Name	Lysine-specific demethylase 5A (EC 1.14.11.-) (Histone demethylase JARID1A) (Jumonji/ARID domain-containing protein 1A) (Retinoblastoma-binding protein 2) (RBBP-2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 720-800
Specificity	KDM5A Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	185kD
Cell Pathway	Nucleus, nucleolus . Nucleus . Occupies promoters of genes involved in RNA metabolism and mitochondrial function. .
Tissue Specificity	Epithelium, Eye, Testis,
Function	cofactor:Fe(2+).,function:Histone demethylase that specifically demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9', H3 'Lys-27', H3 'Lys-36', H3 'Lys-79' or H4 'Lys-20'. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-4'. May stimulate transcription mediated by nuclear receptors. May be involved in transcriptional regulation of Hox proteins during cell differentiation. May participate in transcriptional repression of cytokines such as CXCL12.,similarity:Belongs to the JARID1 histone demethylase family.,similarity:Contains 1 ARID domain.,similarity:Contains 1 JmjC domain.,similarity:Contains 1 JmjN domain.,similarity:Contains 3 PHD-type zinc

Nanjing BYabscience technology Co.,Ltd



fingers.,subunit:Interacts with the viral protein-binding domain of RB1. Interacts with ESR1, MYC, MYCN and LMO2.,

Background

This gene encodes a member of the Jumonji, AT-rich interactive domain 1 (JARID1) histone demethylase protein family. The encoded protein plays a role in gene regulation through the histone code by specifically demethylating lysine 4 of histone H3. The encoded protein interacts with many other proteins, including retinoblastoma protein, and is implicated in the transcriptional regulation of Hox genes and cytokines. This gene may play a role in tumor progression. [provided by RefSeq, Aug 2013],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658