



DPOG1 Polyclonal Antibody

Catalog No	BYab-05521
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	POLG MDP1 POLG1 POLGA
Protein Name	DNA polymerase subunit gamma-1 (EC 2.7.7.7) (Mitochondrial DNA polymerase catalytic subunit) (PolG-alpha)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DPOG1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	136kD
Cell Pathway	Mitochondrion . Mitochondrion matrix, mitochondrion nucleoid .
Tissue Specificity	Brain,Lymph,Testis,
Function	catalytic activity:Deoxynucleoside triphosphate + DNA(n) = diphosphate + DNA(n+1).,cofactor:Magnesium.,disease:Defects in POLG are a cause of Alpers-Huttenlocher syndrome (AHS) [MIM:203700]; also called Alpers diffuse degeneration of cerebral gray matter with hepatic cirrhosis. AHS is an autosomal recessive hepatocerebral syndrome. The typical course of AHS includes severe developmental delay, intractable seizures, liver failure, and death in childhood. Refractory seizures, cortical blindness, progressive liver dysfunction, and acute liver failure after exposure to valproic acid are considered diagnostic features. The neuropathological hallmarks of AHS are neuronal loss, spongiform degeneration, and astrogliosis of the visual cortex. Liver biopsy results show steatosis, often progressing to cirrhosis.,disease:Defects in POLG are a cause of mitochondrial neurogastrointestinal encephalopat

Nanjing BYabscience technology Co.,Ltd

**Background**

Mitochondrial DNA polymerase is heterotrimeric, consisting of a homodimer of accessory subunits plus a catalytic subunit. The protein encoded by this gene is the catalytic subunit of mitochondrial DNA polymerase. The encoded protein contains a polyglutamine tract near its N-terminus that may be polymorphic. Defects in this gene are a cause of progressive external ophthalmoplegia with mitochondrial DNA deletions 1 (PEOA1), sensory ataxic neuropathy dysarthria and ophthalmoparesis (SANDO), Alpers-Huttenlocher syndrome (AHS), and mitochondrial neurogastrointestinal encephalopathy syndrome (MNGIE). Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658