



NEIL2 Polyclonal Antibody

Catalog No	BYab-05573
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	NEIL2
Protein Name	Endonuclease 8-like 2 (EC 3.2.2.-) (EC 4.2.99.18) (DNA glycosylase/AP lyase Neil2) (DNA-(apurinic or apyrimidinic site) lyase Neil2) (Endonuclease VIII-like 2) (Nei homolog 2) (NEH2) (Nei-like protein)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	NEIL2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	36kD
Cell Pathway	Nucleus .
Tissue Specificity	Detected in testis, skeletal muscle, heart, brain, placenta, lung, pancreas, kidney and liver.
Function	catalytic activity:Removes damaged bases from DNA, leaving an abasic site.,catalytic activity:The C-O-P bond 3' to the apurinic or apyrimidinic site in DNA is broken by a beta-elimination reaction, leaving a 3'-terminal unsaturated sugar and a product with a terminal 5'-phosphate.,domain:The zinc-finger domain is important for DNA binding.,enzyme regulation:Acetylation of Lys-50 leads to loss of DNA nicking activity. Acetylation of Lys-154 has no effect.,function:Involved in base excision repair of DNA damaged by oxidation or by mutagenic agents. Has DNA glycosylase activity towards 5-hydroxyuracil and other oxidized derivatives of cytosine with a preference for mismatched double stranded DNA (DNA bubbles). Has low or no DNA glycosylase activity towards thymine glycol,

Nanjing BYabscience technology Co.,Ltd



2-hydroxyadenine, hypoxanthine and 8-oxoguanine. Has AP (apurinic/aprimidinic) lyase activity and introduces nicks in t

Background

NEIL2 belongs to a class of DNA glycosylases homologous to the bacterial Fpg/Nei family. These glycosylases initiate the first step in base excision repair by cleaving bases damaged by reactive oxygen species and introducing a DNA strand break via the associated lyase reaction (Bandaru et al., 2002 [PubMed 12509226])[supplied by OMIM, Mar 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658