



NOG2 Polyclonal Antibody

Catalog No	BYab-05879
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	GNL2 NGP1
Protein Name	Nucleolar GTP-binding protein 2 (Autoantigen NGP-1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 670-750
Specificity	NOG2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	80kD
Cell Pathway	Nucleus, nucleolus .
Tissue Specificity	Widely expressed, with the highest expression level in testis.
Function	function:GTPase that associates with pre-60S ribosomal subunits in the nucleolus and is required for their nuclear export and maturation.,similarity:Belongs to the MMR1/HSR1 GTP-binding protein family. NOG2 subfamily.,tissue specificity:Ubiquitously expressed at relatively low levels in all human tissues tested, with the highest level of expression in the testes.,
Background	function:GTPase that associates with pre-60S ribosomal subunits in the nucleolus and is required for their nuclear export and maturation.,similarity:Belongs to the MMR1/HSR1 GTP-binding protein family. NOG2 subfamily.,tissue specificity:Ubiquitously expressed at relatively low levels in all human tissues tested, with the highest level of expression in the testes.,

Nanjing BYabscience technology Co.,Ltd

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images
