



PPIG Polyclonal Antibody

Catalog No	BYab-05902
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	PPIG
Protein Name	Peptidyl-prolyl cis-trans isomerase G (PPlase G) (Peptidyl-prolyl isomerase G) (EC 5.2.1.8) (CASP10) (Clk-associating RS-cyclophilin) (CARS-Cyp) (CARS-cyclophilin) (SR-cyclophilin) (SR-cyp) (SRcyp) (C
Immunogen	Synthesized peptide derived from human protein . at AA range: 290-370
Specificity	PPIG Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	82kD
Cell Pathway	Nucleus matrix . Nucleus speckle . Colocalizes with RNA splicing factors at nuclear speckles. .
Tissue Specificity	Ubiquitous.
Function	catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0).,domain:The RS domain is required for the interaction with the phosphorylated C-terminal domain of RNA polymerase II.,enzyme regulation:Cyclosporin A (CsA)-sensitive.,function:PPLases accelerate the folding of proteins.,function:PPLases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.,function:PPLases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. May be implicated in the folding, transport, and assembly of proteins. May play an important role in the regulation of pre-mRNA splicing.,PTM:Phosphorylated upon

Nanjing BYabscience technology Co.,Ltd



DNA damage, probably by ATM or ATR.,similarity:Belongs to the cyclophilin-type PPlase family.,similarity:Contains 1 PPlase cyclophilin-type domain.,subcell

Background

catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0).,domain:The RS domain is required for the interaction with the phosphorylated C-terminal domain of RNA polymerase II.,enzyme regulation:Cyclosporin A (CsA)-sensitive.,function:PPlases accelerate the folding of proteins.,function:PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.,function:PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. May be implicated in the folding, transport, and assembly of proteins. May play an important role in the regulation of pre-mRNA splicing.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the cyclophilin-type PPlase family.,similarity:Contains 1 PPlase cyclophilin-type domain.,subcellular location:Colocalizes with RNA splicing factors at nuclear speckles.,subunit:Interacts with CLK1, PNN and with the phosphorylated C-terminal domain of RNA polymerase II.,tissue specificity:Ubiquitous.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658