



PR40A Polyclonal Antibody

Catalog No	BYab-05969
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	PRPF40A FBP11 FLAF1 FNBP3 HIP10 HYPA HSPC225
Protein Name	Pre-mRNA-processing factor 40 homolog A (Fas ligand-associated factor 1) (Formin-binding protein 11) (Formin-binding protein 3) (Huntingtin yeast partner A) (Huntingtin-interacting protein 10) (HIP-10)
Immunogen	Synthesized peptide derived from human protein . at AA range: 450-530
Specificity	PR40A Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	105kD
Cell Pathway	Nucleus speckle . Nucleus matrix . Colocalizes with AKAP8L in the nuclear matrix. .
Tissue Specificity	Expressed in the brain cortex (at protein level). Widely expressed.
Function	domain:The WW domains are essential for localization to nuclear speckles.,function:Binds to WASL/N-WASP and suppresses its translocation from the nucleus to the cytoplasm, thereby inhibiting its cytoplasmic function (By similarity). May be involved in pre-mRNA splicing.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 409.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the PRPF40 family.,similarity:Contains 2 WW domains.,similarity:Contains 5 FF domains.,subunit:Interacts with the N-terminus of HTT. Interacts with the phosphorylated carboxyterminal domain of POLR2A. Interacts with SF1, SRPK1 CARD8, ATBF1 and MECP2 (By similarity). Interacts

Nanjing BYabscience technology Co.,Ltd



through the WW domains with formin proline-rich regions and with WASL/N-WASP.,tissue specificity:Widely expressed.,

Background

domain:The WW domains are essential for localization to nuclear speckles.,function:Binds to WASL/N-WASP and suppresses its translocation from the nucleus to the cytoplasm, thereby inhibiting its cytoplasmic function (By similarity). May be involved in pre-mRNA splicing.,sequence caution:Contaminating sequence. Potential poly-A sequence starting in position 409.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the PRPF40 family.,similarity:Contains 2 WW domains.,similarity:Contains 5 FF domains.,subunit:Interacts with the N-terminus of HTT. Interacts with the phosphorylated carboxyterminal domain of POLR2A. Interacts with SF1, SRPK1 CARD8, ATBF1 and MECP2 (By similarity). Interacts through the WW domains with formin proline-rich regions and with WASL/N-WASP.,tissue specificity:Widely expressed.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658