



CROCC Polyclonal Antibody

Catalog No	BYab-06115
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CROCC KIAA0445
Protein Name	Rootletin (Ciliary rootlet coiled-coil protein)
Immunogen	Synthesized peptide derived from human protein . at AA range: 220-300
Specificity	CROCC Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	221kD
Cell Pathway	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole . Cytoplasm, cytoskeleton, cilium basal body . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . In ciliated cells, associated with ciliary rootlets. In non-ciliated cells, localized between, around and at the proximal ends of the centrioles. Dissociates from the centrioles at the onset of mitosis and reassociates with them at anaphase. .
Tissue Specificity	Brain,Medulloblastoma,
Function	function:Major structural component of the ciliary rootlet, a cytoskeletal-like structure in ciliated cells which originates from the basal body at the proximal end of a cilium and extends proximally toward the cell nucleus. Contributes to centrosome cohesion before mitosis.,PTM:Phosphorylated by NEK2 which may regulate its association with centrosomes. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the rootletin family.,subcellular location:In ciliated cells, associated with ciliary rootlets. In non-ciliated cells, localized between, around and at the proximal ends of the centrioles. Dissociates

Nanjing BYabscience technology Co.,Ltd



	from the centrioles at the onset of mitosis and reassociates with them at anaphase.,subunit:Homopolymer. Interacts with KLC3, NEK2 and the N-terminus of CEP250.,
Background	function:Major structural component of the ciliary rootlet, a cytoskeletal-like structure in ciliated cells which originates from the basal body at the proximal end of a cilium and extends proximally toward the cell nucleus. Contributes to centrosome cohesion before mitosis.,PTM:Phosphorylated by NEK2 which may regulate its association with centrosomes. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the rootletin family.,subcellular location:In ciliated cells, associated with ciliary rootlets. In non-ciliated cells, localized between, around and at the proximal ends of the centrioles. Dissociates from the centrioles at the onset of mitosis and reassociates with them at anaphase.,subunit:Homopolymer. Interacts with KLC3, NEK2 and the N-terminus of CEP250.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658