



SRPK2 Polyclonal Antibody

Catalog No	BYab-06140
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	SRPK2
Protein Name	SRSF protein kinase 2 (EC 2.7.11.1) (SFRS protein kinase 2) (Serine/arginine-rich protein-specific kinase 2) (SR-protein-specific kinase 2) [Cleaved into: SRSF protein kinase 2 N-terminal; SRSF protei
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SRPK2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	75kD
Cell Pathway	Cytoplasm . Nucleus, nucleoplasm . Nucleus speckle . Chromosome . Shuttles between the nucleus and the cytoplasm (PubMed:19592491, PubMed:21157427, PubMed:21056976). KAT5/TIP60 inhibits its nuclear translocation (PubMed:21157427). Phosphorylation at Thr-492 by PKB/AKT1 promotes nuclear translocation (PubMed:19592491). Preferentially localizes across the entire gene coding region (PubMed:28076779). During transcription, accumulates at chromatin loci where unscheduled R-loops form and colocalizes with paused 'Ser-5'-phosphorlyated POLR2A/RNA polymerase II and helicase DDX23 (PubMed:28076779). .
Tissue Specificity	Highly expressed in brain, moderately expressed in heart and skeletal muscle and at low levels in lung, liver, and kidney.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Ser-52 and Ser-588.,function:Phosphorylates RS

Nanjing BYabscience technology Co.,Ltd



domain-containing proteins, such as SFRS1 and SFRS2 on serine residues. Role in spliceosome assembly and in mediating the trafficking of splicing factors. Appears to mediate HBV core protein phosphorylation which is a prerequisite for pregenomic RNA encapsidation into viral capsids.,sequence caution:The cDNA appears to contain a duplicated region.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,tissue specificity:Highly expressed in brain, moderately expressed in heart and skeletal muscle and at low levels in lung, liver, and kidney.,

Background

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Ser-52 and Ser-588.,function:Phosphorylates RS domain-containing proteins, such as SFRS1 and SFRS2 on serine residues. Role in spliceosome assembly and in mediating the trafficking of splicing factors. Appears to mediate HBV core protein phosphorylation which is a prerequisite for pregenomic RNA encapsidation into viral capsids.,sequence caution:The cDNA appears to contain a duplicated region.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,tissue specificity:Highly expressed in brain, moderately expressed in heart and skeletal muscle and at low levels in lung, liver, and kidney.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658