



TRPM7 Polyclonal Antibody

Catalog No	BYab-06313
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	TRPM7 CHAK1 LTRPC7
Protein Name	Transient receptor potential cation channel subfamily M member 7 (EC 2.7.11.1) (Channel-kinase 1) (Long transient receptor potential channel 7) (LTrpC-7) (LTrpC7)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TRPM7 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	205kD
Cell Pathway	Membrane ; Multi-pass membrane protein .
Tissue Specificity	Colon,Liver,Placenta,
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Binds 1 zinc ion per subunit.,disease:Defects in TRPM7 influence susceptibility to amyotrophic lateral sclerosis-parkinsonism/dementia complex type 1 [MIM:105500]; also called amyotrophic lateral sclerosis-parkinsonism/dementia complex of Guam or Guam disease. Amyotrophic lateral sclerosis-parkinsonism/dementia complex type 1 is a neurodegenerative disorder with unusually high incidence among the Chamorro people of the Western Pacific Islands of Guam. Both amyotrophic lateral sclerosis and parkinsonism-dementia are chronic, progressive, and uniformly fatal disorders in this population. Both diseases are known to occur in the same kindred, the same sibship, and even the same individual.,function:Essential ion channel and serine/threonine-protein kinase. Divalent cation channel permeable


Nanjing BYabscience technology Co.,Ltd



to calcium and magnesium. Has a c

Background	transient receptor potential cation channel subfamily M member 7 (TRPM7) Homo sapiens The protein encoded by this gene is both an ion channel and a serine/threonine protein kinase. The kinase activity is essential for the ion channel function, which serves to increase intracellular calcium levels and to help regulate magnesium ion homeostasis. Defects in this gene are a cause of amyotrophic lateral sclerosis-parkinsonism/dementia complex of Guam. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of K562 and A549 cell lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658