



TNNT3 Polyclonal Antibody

Catalog No	BYab-06323
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	TNNT3
Protein Name	Troponin T, fast skeletal muscle (TnTf) (Beta-TnTF) (Fast skeletal muscle troponin T) (fTnT)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TNNT3 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	29kD
Cell Pathway	cytosol,troponin complex,
Tissue Specificity	In fetal and adult fast skeletal muscles, with a higher level expression in fetal than in adult muscle.
Function	alternative products:Additional isoforms seem to exist,disease:Defects in TNNT3 are a cause of distal arthrogryposis type 2B (DA2B) [MIM:601680]; also known as arthrogryposis multiplex congenita, distal, type 2B (AMCD2B). DA2B is a form of inherited multiple congenital contractures. Affected individuals have vertical talus, ulnar deviation in the hands, severe camptodactyly, and a distinctive face characterized by a triangular shape, prominent nasolabial folds, small mouth and a prominent chin.,function:Troponin T is the tropomyosin-binding subunit of troponin, the thin filament regulatory complex which confers calcium-sensitivity to striated muscle actomyosin ATPase activity.,similarity:Belongs to the troponin T family.,tissue specificity:In fetal and adult fast skeletal muscles, with a higher level expression in fetal than in adult muscle.,

Nanjing BYabscience technology Co.,Ltd



Background

The binding of Ca^{2+} to the trimeric troponin complex initiates the process of muscle contraction. Increased Ca^{2+} concentrations produce a conformational change in the troponin complex that is transmitted to tropomyosin dimers situated along actin filaments. The altered conformation permits increased interaction between a myosin head and an actin filament which, ultimately, produces a muscle contraction. The troponin complex has protein subunits C, I, and T. Subunit C binds Ca^{2+} and subunit I binds to actin and inhibits actin-myosin interaction. Subunit T binds the troponin complex to the tropomyosin complex and is also required for Ca^{2+} -mediated activation of actomyosin ATPase activity. There are 3 different troponin T genes that encode tissue-specific isoforms of subunit T for fast skeletal-, slow skeletal-, and cardiac-muscle. This gene encodes fast skeletal troponin T protein; als


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night