



SNUT1 Polyclonal Antibody

Catalog No	BYab-06330
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	SART1
Protein Name	U4/U6.U5 tri-snRNP-associated protein 1 (SNU66 homolog) (hSnu66) (Squamous cell carcinoma antigen recognized by T-cells 1) (SART-1) (hSART-1) (U4/U6.U5 tri-snRNP-associated 110 kDa protein) (allergen)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SNUT1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	88kD
Cell Pathway	Nucleus . Found in the nucleus of mitogen-activated peripheral blood mononuclear cells (PBMCs), tumor cells, or normal cell lines, but not in normal tissues except testis and fetal liver or in unstimulated PBMCs, suggesting preferential expression in proliferating cells.
Tissue Specificity	Ubiquitously expressed.
Function	allergen:Causes an allergic reaction in human. Binds to IgE from atopic dermatitis (AD) patients. Identified as an IgE autoantigen in atopic dermatitis (AD) patients with severe skin manifestations.,alternative products:A number of isoforms may be produced,function:May play a role in mRNA splicing. May also bind to DNA.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Sumoylated by SUMO2.,similarity:Belongs to the SNU66/SART1 family.,subcellular location:Found in the nucleus of mitogen-activated peripheral blood mononuclear cells (PBMCs), tumor cells, or normal cell lines, but not in

Nanjing BYabscience technology Co.,Ltd



normal tissues except testis and fetal liver or in unstimulated PBMCs, suggesting preferential expression in proliferating cells.,subunit:Identified in the spliceosome C complex, at least composed of AQR, ASCC3L1, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38,

Background

This gene encodes two proteins, the SART1(800) protein expressed in the nucleus of the majority of proliferating cells, and the SART1(259) protein expressed in the cytosol of epithelial cancers. The SART1(259) protein is translated by the mechanism of -1 frameshifting during posttranscriptional regulation; its full-length sequence is not published yet. The two encoded proteins are thought to be involved in the regulation of proliferation. Both proteins have tumor-rejection antigens. The SART1(259) protein possesses tumor epitopes capable of inducing HLA-A2402-restricted cytotoxic T lymphocytes in cancer patients. This SART1(259) antigen may be useful in specific immunotherapy for cancer patients and may serve as a paradigmatic tool for the diagnosis and treatment of patients with atopy. The SART1(259) protein is found to be essential for the recruitment of the tri-snR

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658