



UBA6 Polyclonal Antibody

Catalog No	BYab-06353
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	UBA6 MOP4 UBE1L2
Protein Name	Ubiquitin-like modifier-activating enzyme 6 (Ubiquitin-activating enzyme 6) (Monocyte protein 4) (MOP-4) (Ubiquitin-activating enzyme E1-like protein 2) (E1-L2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	UBA6 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	115kD
Cell Pathway	cytoplasm,cytosol,integral component of membrane,
Tissue Specificity	Widely expressed. Isoform 2 is predominantly expressed in testis with higher expression in adult testis than in fetal testis.
Function	alternative products:Named isoforms=3,function:Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding an ubiquitin-E1 thioester and free AMP. Specific for ubiquitin, does not activate ubiquitin-like peptides. Differs from UBE1 in its specificity for substrate E2 charging. Does not charge cell cycle E2s, such as CDC34. Isoform 2 may play a key role in ubiquitin system and may influence spermatogenesis and male fertility.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the ubiquitin-activating E1 family.,tissue specificity:Widely expressed. Isoform 2 is predominantly expressed in testis with higher expression in adult testis than in fetal testis.,

Nanjing BYabscience technology Co.,Ltd



Background	Modification of proteins with ubiquitin (UBB; MIM 191339) or ubiquitin-like proteins controls many signaling networks and requires a ubiquitin-activating enzyme (E1), a ubiquitin conjugating enzyme (E2), and a ubiquitin protein ligase (E3). UBE1L2 is an E1 enzyme that initiates the activation and conjugation of ubiquitin-like proteins (Jin et al., 2007 [PubMed 17597759]).[supplied by OMIM, Mar 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images