



KDM4D Polyclonal Antibody

Catalog No	BYab-06510
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	KDM4D JHDM3D JMJD2D
Protein Name	Lysine-specific demethylase 4D (EC 1.14.11.-) (JmjC domain-containing histone demethylation protein 3D) (Jumonji domain-containing protein 2D)
Immunogen	Synthesized peptide derived from human protein . at AA range: 400-480
Specificity	KDM4D Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	57kD
Cell Pathway	Nucleus .
Tissue Specificity	Brain,Embryo,Teratocarcinoma,Testis,
Function	caution:It is uncertain whether Met-1 or Met-4 is the initiator.,cofactor:Binds 1 Fe(2+) ion per subunit.,function:Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20'. Demethylates both di- and trimethylated H3 'Lys-9' residue, while it has no activity on monomethylated residues. Demethylation of Lys residue generates formaldehyde and succinate.,similarity:Belongs to the JHDM3 histone demethylase family.,similarity:Contains 1 JmjC domain.,similarity:Contains 1 JmjN domain.,
Background	caution:It is uncertain whether Met-1 or Met-4 is the initiator.,cofactor:Binds 1 Fe(2+) ion per subunit.,function:Histone demethylase that specifically

Nanjing BYabscience technology Co.,Ltd



demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20'. Demethylates both di- and trimethylated H3 'Lys-9' residue, while it has no activity on monomethylated residues. Demethylation of Lys residue generates formaldehyde and succinate.,similarity:Belongs to the JHDM3 histone demethylase family.,similarity:Contains 1 JmjC domain.,similarity:Contains 1 JmjN domain.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658