



DIRC2 Polyclonal Antibody

Catalog No	BYab-06555
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	DIRC2
Protein Name	Disrupted in renal carcinoma protein 2 (Disrupted in renal cancer protein 2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 170-250
Specificity	DIRC2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	52kD
Cell Pathway	Lysosome membrane ; Multi-pass membrane protein .
Tissue Specificity	Ubiquitous. Expressed in proximal tubular cells of the kidney.
Function	disease:A chromosomal aberration involving DIRC2 is found in familial renal cell carcinoma 1 (RCC1) [MIM:144700]. Translocation t(2;3)(q35;q21).,similarity:Belongs to the major facilitator superfamily.,tissue specificity:Ubiquitous. Expressed in proximal tubular cells of the kidney.,
Background	This gene encodes a membrane-bound protein from the major facilitator superfamily of transporters. Disruption of this gene by translocation has been associated with haplo-insufficiency and renal cell carcinomas. Alternatively spliced transcript variants have been described, but their biological validity has not yet been determined. [provided by RefSeq, Jul 2008],

Nanjing BYabscience technology Co.,Ltd



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images