



I12R2 Polyclonal Antibody

Catalog No	BYab-06621
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	IL12RB2
Protein Name	Interleukin-12 receptor subunit beta-2 (IL-12 receptor subunit beta-2) (IL-12R subunit beta-2) (IL-12R-beta-2) (IL-12RB2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 270-350
Specificity	I12R2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	94kD
Cell Pathway	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Isoform 2 is expressed at similar levels in both naive and activated T-cells.
Function	developmental stage:Maximum levels in Th1 cells between day 3 and day 8 of activation.,domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:Receptor for interleukin-12. This subunit is the signaling component coupling to the JAK2/STAT4 pathway. Promotes the proliferation of T-cells as well as NK cells. Induces the promotion of T-cells towards the Th1 phenotype by strongly enhancing IFN-gamma production.,induction:In vitro, up-regulated by interferon alpha.,polymorphism:Heterozygotic variants Gly-313 and Arg-720 are associated with atopy, an immunological condition that can lead to clinical symptoms such as allergic rhinitis, sinusitis, asthma and eczema.,PTM:On IL12 binding, phosphorylated on C-terminal ty

Nanjing BYabscience technology Co.,Ltd

**Background**

interleukin 12 receptor subunit beta 2(IL12RB2) Homo sapiens The protein encoded by this gene is a type I transmembrane protein identified as a subunit of the interleukin 12 receptor complex. The coexpression of this and IL12RB1 proteins was shown to lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent signaling. The expression of this gene is up-regulated by interferon gamma in Th1 cells, and plays a role in Th1 cell differentiation. The up-regulation of this gene is found to be associated with a number of infectious diseases, such as Crohn's disease and leprosy, which is thought to contribute to the inflammatory response and host defense. Several transcript variants encoding different isoforms and non-protein coding transcripts have been found for this gene. [provided by RefSeq, Apr 2012],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658