



OLIG1 Polyclonal Antibody

Catalog No	BYab-07032
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	OLIG1 BHLHB6 BHLHE21
Protein Name	Oligodendrocyte transcription factor 1 (Oligo1) (Class B basic helix-loop-helix protein 6) (bHLHb6) (Class E basic helix-loop-helix protein 21) (bHLHe21)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	OLIG1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	29kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas is highly variable.
Function	function:Promotes formation and maturation of oligodendrocytes, especially within the brain. Cooperates with OLIG2 to establish the pMN domain of the embryonic neural tube.,induction:By SHH.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,tissue specificity:Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas is highly variable.,
Background	function:Promotes formation and maturation of oligodendrocytes, especially within the brain. Cooperates with OLIG2 to establish the pMN domain of the embryonic neural tube.,induction:By SHH.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,tissue specificity:Expressed in the brain, in oligodendrocytes. Strongly

Nanjing BYabscience technology Co.,Ltd



expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas is highly variable.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4°over night