



AMGO2 Polyclonal Antibody

Catalog No	BYab-07231
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	AMIGO2 ALI1
Protein Name	Amphoterin-induced protein 2 (AMIGO-2) (Alivin-1) (Differentially expressed in gastric adenocarcinomas) (DEGA)
Immunogen	Synthesized peptide derived from human protein . at AA range: 270-350
Specificity	AMGO2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	57kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein . Nucleus . Associated with nucleus as well as plasma membrane. Restricted to somata of cerebellar as well as hippocampal neurons (By similarity). .
Tissue Specificity	Highest levels in breast, ovary, cervix, and uterus. Lower levels in lung, colon, and rectum. Differentially expressed in 56% of thyroid, 57% of pancreatic and 45% of stomach cancers.
Function	function:Required for depolarization-dependent survival of cultured cerebellar granule neurons. May mediate homophilic as well as heterophilic cell-cell interaction with AMIGO1 or AMIGO3. May contribute to signal transduction through its intracellular domain. May be required for tumorigenesis of a subset of gastric adenocarcinomas.,similarity:Belongs to the immunoglobulin superfamily. AMIGO family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 6 LRR (leucine-rich) repeats.,subcellular location:Associated with nucleus as well as plasma membrane. Restricted to somata of cerebellar as well as hippocampal neurons.,subunit:Binds itself as well

Nanjing BYabscience technology Co.,Ltd



	as AMIGO1 and AMIGO3.,tissue specificity:Highest levels in breast, ovary, cervix, and uterus. Lower levels in lung, colon, and rectum. Differentially expressed in 56% of thyroid, 57% of pancreatic and 45% of sto
Background	function:Required for depolarization-dependent survival of cultured cerebellar granule neurons. May mediate homophilic as well as heterophilic cell-cell interaction with AMIGO1 or AMIGO3. May contribute to signal transduction through its intracellular domain. May be required for tumorigenesis of a subset of gastric adenocarcinomas.,similarity:Belongs to the immunoglobulin superfamily. AMIGO family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 6 LRR (leucine-rich) repeats.,subcellular location:Associated with nucleus as well as plasma membrane. Restricted to somata of cerebellar as well as hippocampal neurons.,subunit:Binds itself as well as AMIGO1 and AMIGO3.,tissue specificity:Highest levels in breast, ovary, cervix, and uterus. Lower levels in lung, colon, and rectum. Differentially expressed in 56% of thyroid, 57% of pancreatic and 45% of stomach cancers.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658