



JPH2 Polyclonal Antibody

Catalog No	BYab-07318
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	JPH2 JP2
Protein Name	Junctophilin-2 (JP-2) (Junctophilin type 2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 611-660
Specificity	JPH2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	76kD
Cell Pathway	[Junctophilin-2]: Cell membrane ; Peripheral membrane protein . Sarcoplasmic reticulum membrane ; Single-pass type IV membrane protein . Endoplasmic reticulum membrane ; Single-pass type IV membrane protein . The transmembrane domain is anchored in sarcoplasmic reticulum membrane, while the N-terminal part associates with the plasma membrane. In heart cells, it predominantly associates along Z lines within myocytes. In skeletal muscle, it is specifically localized at the junction of A and I bands. . ; [Junctophilin-2 N-terminal fragment]: Nucleus . Accumulates in the nucleus of stressed hearts. .
Tissue Specificity	Specifically expressed in skeletal muscle and heart.
Function	domain:The MORN (membrane occupation and recognition nexus) repeats contribute to the plasma membrane binding, possibly by interacting with phospholipids.,function:Contributes to the stabilization of the junctional membrane complexes, which are common to excitable cells and mediate cross-talk between cell surface and intracellular ion channels. Probably acts by anchoring the plasma membrane and endoplasmic/sarcoplasmic reticulum. Contributes to the

Nanjing BYabscience technology Co.,Ltd



construction of skeletal muscle triad junctions, and plays an essential role in heart development.,similarity:Belongs to the junctophilin family.,similarity:Contains 8 MORN repeats.,subcellular location:Localized predominantly on the plasma membrane. The transmembrane domain is anchored in endoplasmic/sarcoplasmic reticulum membrane, while the N-terminal part associates with the plasma membrane. In heart cells, it predominantly associates alone

Background

Junctional complexes between the plasma membrane and endoplasmic/sarcoplasmic reticulum are a common feature of all excitable cell types and mediate cross talk between cell surface and intracellular ion channels. The protein encoded by this gene is a component of junctional complexes and is composed of a C-terminal hydrophobic segment spanning the endoplasmic/sarcoplasmic reticulum membrane and a remaining cytoplasmic domain that shows specific affinity for the plasma membrane. This gene is a member of the junctophilin gene family. Alternative splicing has been observed at this locus and two variants encoding distinct isoforms are described. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658