



TSHR Polyclonal Antibody

Catalog No	BYab-07557
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	TSHR LGR3
Protein Name	Thyrotropin receptor (Thyroid-stimulating hormone receptor) (TSH-R)
Immunogen	Synthesized peptide derived from human protein . at AA range: 280-360
Specificity	TSHR Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	84kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Basolateral cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Expressed in thyroide cells (at protein level) (PubMed:11847099). Expressed in the thyroid (PubMed:2610690).
Function	alternative products:Additional isoforms seem to exist,disease:Autoantibodies directed against the TSH receptor are directly responsible for the pathogenesis and hyperthyroidism of Graves disease (GRD) [MIM:275000]. Antibody interaction with the TSH receptor results in an uncontrolled receptor stimulation.,disease:Defects in TSHR are a cause of hyperthyroidism [MIM:603372]. Various types are known: autosomal dominant non-autoimmune hyperthyroidism (ADNH); sporadic congenital hyperthyroidism (SCH); hyperthyroidism associated with autonomously functioning thyroid nodules (AFTN), toxic multinodular goiter (TMNG) and hyperfunctioning thyroid adenomas (HTA). TMNG encompasses a spectrum of different clinical entities, ranging from a single hyperfunctioning nodule within an enlarged thyroid, to multiple

Nanjing BYabscience technology Co.,Ltd



hyperfunctioning areas scattered throughout the gland. HTA are discrete encapsulated neoplas

Background

The protein encoded by this gene is a membrane protein and a major controller of thyroid cell metabolism. The encoded protein is a receptor for thyrothropin and thyrostimulin, and its activity is mediated by adenylate cyclase. Defects in this gene are a cause of several types of hyperthyroidism. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images