



WDR20 rabbit pAb

Catalog No BYab-08059 Isotype IgG Reactivity Human; Mouse Applications WB Gene Name WDR20 Protein Name WDR20 Immunogen Synthesized peptide derived from human WDR20 AA range: 282-332 Specificity This antibody detects endogenous levels of WDR20 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma, Testis, Function similarity:Contains 5 WD repeats. Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubjequitinating enzyme complex. Multi		
Reactivity Applications WB Gene Name WDR20 Protein Name Immunogen Synthesized peptide derived from human WDR20 AA range: 282-332 Specificity This antibody detects endogenous levels of WDR20 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Armygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis, Function similarity:Contains 5 WD repeats. Background Tisgene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Catalog No	BYab-08059
Applications WB Gene Name WDR20 Protein Name WDR20 Immunogen Synthesized peptide derived from human WDR20 AA range: 282-332 Specificity This antibody detects endogenous levels of WDR20 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats. Background This gene encodes w WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], Matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name WDR20 Protein Name WDR20 Immunogen Synthesized peptide derived from human WDR20 AA range: 282-332 Specificity This antibody detects endogenous levels of WDR20 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats. Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], Matters needing Avoid repeated freezing and thawing!	Reactivity	Human; Mouse
Protein Name WDR20 Immunogen Synthesized peptide derived from human WDR20 AA range: 282-332 Specificity This antibody detects endogenous levels of WDR20 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats. Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Applications	WB
Immunogen Synthesized peptide derived from human WDR20 AA range: 282-332 Specificity This antibody detects endogenous levels of WDR20 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Gene Name	WDR20
Specificity This antibody detects endogenous levels of WDR20 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Protein Name	WDR20
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human WDR20 AA range: 282-332
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex, Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of WDR20 at Human/Mouse
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.174% sodium azide.
using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis, Function similarity:Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function similarity: Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis, Function similarity:Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis, Function similarity:Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms WD repeat-containing protein 20 (Protein DMR) Observed Band 60kD Cell Pathway Tissue Specificity Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis, Function similarity:Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Tissue Specificity Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis, Function Similarity:Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Tissue Specificity Amygdala, Brain, Endometrium carcinoma cell line, Epithelium, Neuroblastoma, Testis, Function Similarity: Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Synonyms	WD repeat-containing protein 20 (Protein DMR)
Tissue Specificity Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis, similarity:Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Observed Band	60kD
line,Epithelium,Neuroblastoma,Testis, Function similarity:Contains 5 WD repeats., Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Cell Pathway	
Background This gene encodes a WD repeat-containing protein that functions to preserve and regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Amygdala,Brain,Endometrium carcinoma cell line,Epithelium,Neuroblastoma,Testis,
regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2011], matters needing Avoid repeated freezing and thawing!	Function	similarity:Contains 5 WD repeats.,
	Background	regulate the activity of the USP12-UAF1 deubiquitinating enzyme complex. Multiple alternatively spliced transcript variants have been found for this gene.
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from 3T3 cells, primary antibody was diluted at 1:1000, 4°over night

Nanjing BYabscience technology Co.,Ltd