



OXDD rabbit pAb

Catalog No	BYab-08560
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	DDO
Protein Name	OXDD
Immunogen	Synthesized peptide derived from human OXDD AA range: 17-67
Specificity	This antibody detects endogenous levels of OXDD at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Peroxisome.
Tissue Specificity	
Function	catalytic activity:D-aspartate + H(2)O + O(2) = oxaloacetate + NH(3) + H(2)O(2).,cofactor:FAD or 6-hydroxyflavin adenine dinucleotide.,function:Selectively catalyzes the oxidative deamination of D-aspartate and its N-methylated derivative, N-methyl D-aspartate.,similarity:Belongs to the DAMOX/DASOX family.,
Background	The protein encoded by this gene is a peroxisomal flavoprotein that catalyzes the oxidative deamination of D-aspartate and N-methyl D-aspartate. Flavin adenine dinucleotide or 6-hydroxyflavin adenine dinucleotide can serve as the cofactor in this reaction. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from THP-1 cells, primary antibody was diluted at 1:1000, 4° over night