



# MORN2 rabbit pAb

|                           |                                                                                                                                                                                                                       |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog No                | BYab-08589                                                                                                                                                                                                            |
| Isotype                   | IgG                                                                                                                                                                                                                   |
| Reactivity                | Human; Mouse                                                                                                                                                                                                          |
| Applications              | IHC;IF                                                                                                                                                                                                                |
| Gene Name                 | MORN2 MOPT                                                                                                                                                                                                            |
| Protein Name              | MORN2                                                                                                                                                                                                                 |
| Immunogen                 | Synthesized peptide derived from human MORN2 AA range: 25-75                                                                                                                                                          |
| Specificity               | This antibody detects endogenous levels of MORN2 at Human/Mouse                                                                                                                                                       |
| Formulation               | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                               |
| Source                    | Polyclonal, Rabbit,IgG                                                                                                                                                                                                |
| Purification              | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.                                                                                                             |
| Dilution                  | IHC-p 1: 50-200. IF 1:50-200                                                                                                                                                                                          |
| Concentration             | 1 mg/ml                                                                                                                                                                                                               |
| Purity                    | ≥90%                                                                                                                                                                                                                  |
| Storage Stability         | -20°C/1 year                                                                                                                                                                                                          |
| Synonyms                  |                                                                                                                                                                                                                       |
| Observed Band             |                                                                                                                                                                                                                       |
| Cell Pathway              | Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis. . |
| Tissue Specificity        |                                                                                                                                                                                                                       |
| Function                  | similarity:Contains 2 MORN repeats.,                                                                                                                                                                                  |
| Background                |                                                                                                                                                                                                                       |
| matters needing attention | Avoid repeated freezing and thawing!                                                                                                                                                                                  |

Nanjing BYabs science technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998


监督电话: 15950492658



### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).