



CHIC2 rabbit pAb

Catalog No	BYab-08647
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	CHIC2 BTL
Protein Name	CHIC2
Immunogen	Synthesized peptide derived from human CHIC2 AA range: 111-161
Specificity	This antibody detects endogenous levels of CHIC2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cell membrane . Cytoplasmic vesicle . Also present at a Golgi-like vesicular compartment and at scattered vesicles.
Tissue Specificity	
Function	disease:A chromosomal aberration involving CHIC2 is found in a form of acute myeloid leukemia (AML). Translocation t(4;12)(q12;p13) with ETV6.,PTM:Palmitoylation in the CHIC motif is required for membrane association.,similarity:Belongs to the CHIC family.,subcellular location:Also present at a Golgi-like vesicular compartment and at scattered vesicles.,
Background	This gene encodes a member of the CHIC family of proteins. The encoded protein contains a cysteine-rich hydrophobic (CHIC) motif, and is localized to vesicular structures and the plasma membrane. This gene is associated with some cases of acute myeloid leukemia. [provided by RefSeq, Jul 2008],

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night