



WIPI2 Rabbit Polyclonal Antibody

Catalog No	BYab-10267
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF
Gene Name	WIPI2 CGI-50
Protein Name	WD repeat domain phosphoinositide-interacting protein 2 (WIPI-2) (WIPI49-like protein 2)
Immunogen	Recombinant Protein of WIPI2
Specificity	The antibody detects endogenous WIPI2 protein
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC-p 1:50-300. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	WD repeat domain phosphoinositide-interacting protein 2 (WIPI-2;WIPI49-like protein 2)
Observed Band	49kD
Cell Pathway	Preautophagosomal structure membrane ; Peripheral membrane protein ; Cytoplasmic side . Localizes to omegasomes membranes which are endoplasmic reticulum connected structures at the origin of preautophagosomal structures. Enriched at preautophagosomal structure membranes in response to PtdIns3P. .
Tissue Specificity	Ubiquitously expressed (at protein level). Highly expressed in heart, skeletal muscle and pancreas. Expression is down-regulated in pancreatic and in kidney tumors.
Function	similarity:Contains 3 WD repeats.,tissue specificity:Ubiquitously expressed. Highly expressed in heart, skeletal muscle and pancreas. Expression is down-regulated in pancreatic and in kidney tumors.,
Background	WD repeat domain, phosphoinositide interacting 2(WIPI2) Homo sapiens WD40 repeat proteins are key components of many essential biologic functions. They regulate the assembly of multiprotein complexes by presenting a beta-propeller platform for simultaneous and reversible protein-protein

Nanjing BYabscience technology Co.,Ltd



interactions. Members of the WIPI subfamily of WD40 repeat proteins, such as WIPI2, have a 7-bladed propeller structure and contain a conserved motif for interaction with phospholipids (Proikas-Cezanne et al., 2004 [PubMed 15602573]).[supplied by OMIM, Mar 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).