



Hic-5 (Phospho-Tyr60) rabbit pAb


Catalog No	BYab-10506
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	TGFB1I1 ARA55
Protein Name	Hic-5 (Phospho-Tyr60)
Immunogen	Synthesized peptide derived from human Hic-5 (Phospho-Tyr60)
Specificity	This antibody detects endogenous levels of Hic-5 (Phospho-Tyr60) at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.162% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Transforming growth factor beta-1-induced transcript 1 protein (Androgen receptor coactivator 55 kDa protein) (Androgen receptor-associated protein of 55 kDa) (Hydrogen peroxide-inducible clone 5 protein) (Hic-5)
Observed Band	50kD
Cell Pathway	Cell junction, focal adhesion. Nucleus matrix. Cytoplasm, cytoskeleton. Associated with the actin cytoskeleton; colocalizes with stress fibers.
Tissue Specificity	Expressed in platelets, smooth muscle and prostate stromal cells (at protein level).
Function	domain:The LD (leucine and aspartate-rich) motif 3 mediates interaction with GIT1 and functions as a nuclear export signal.,domain:The LIM zinc-binding domains mediate glucocorticoid receptor coactivation and interaction with AR, CRIP2, ILK, LIMS1, NR3C1, PPARG, TCF3, TCF7L2, SLC6A3 and SMAD3. The LIM zinc-binding 2 and LIM zinc-binding 3 domains mediate targeting to focal adhesions and actin stress fibers. The LIM zinc-binding 3 and LIM zinc-binding 4 domains mediate interaction with TRAF4 and MAPK15. The LIM zinc-binding 4 domain mediates interaction with HSPB1, homooligomerization and targeting to the nuclear matrix. The LIM zinc-binding 3 domain mediates interaction with

Nanjing BYabscience technology Co.,Ltd



	PTPN12.,function:Functions as a molecular adapter coordinating multiple protein-protein interactions at the focal adhesion complex and in the nucleus. Links various intracellular signaling modules to plasma membrane
Background	transforming growth factor beta 1 induced transcript 1(TGFB1I1) Homo sapiens This gene encodes a coactivator of the androgen receptor, a transcription factor which is activated by androgen and has a key role in male sexual differentiation. The encoded protein is thought to regulate androgen receptor activity and may have a role to play in the treatment of prostate cancer. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of 1 A431 cell 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000