



MDA5 Phospho-ser828 rabbit pAb

Catalog No	BYab-10568
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	IFIH1 MDA5 RH116
Protein Name	MDA5 ser828
Immunogen	Synthesized peptide derived from human MDA5 ser828
Specificity	This antibody detects endogenous levels of MDA5 ser828 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.224% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Interferon-induced helicase C domain-containing protein 1 (EC 3.6.4.13) (Clinically amyopathic dermatomyositis autoantigen 140 kDa) (CADM-140 autoantigen) (Helicase with 2 CARD domains) (Helicard) (Interferon-induced with helicase C domain protein 1) (Melanoma differentiation-associated protein 5) (MDA-5) (Murabutide down-regulated protein) (RIG-I-like receptor 2) (RLR-2) (RNA helicase-DEAD box protein 116)
Observed Band	
Cell Pathway	Cytoplasm . Nucleus . Mitochondrion . Upon viral RNA stimulation and ISGylation, translocates from cytosol to mitochondrion. May be found in the nucleus, during apoptosis.
Tissue Specificity	Widely expressed, at a low level. Expression is detected at slightly highest levels in placenta, pancreas and spleen and at barely levels in detectable brain, testis and lung.
Function	disease:Genetic variation in IFIH1 is associated with insulin-dependent diabetes mellitus 19 (IDDM19) [MIM:610155].,function:RNA helicase that, through its ATP-dependent unwinding of RNA, may function to promote message

Nanjing BYabscience technology Co.,Ltd



degradation by specific RNases. Seems to have growth suppressive properties. Involved in innate immune defense against viruses. Upon interaction with intracellular dsRNA produced during viral replication, triggers a transduction cascade involving MAVS/IPS1, which results in the activation of NF-kappa-B, IRF3 and IRF7 and the induction of the expression of antiviral cytokines such as IFN-beta and RANTES (CCL5). ATPase activity is specifically induced by dsRNA. Essential for the production of interferons in response to picornaviruses.,induction:By IFN-beta and TNF-alpha.,miscellaneous:In HIV-1 infected HeLa-CD4 cells, overexpression of IFIH1 results in a great increase in t

Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein that is upregulated in response to treatment with beta-interferon and a protein kinase C-activating compound, mezerein. Irreversible reprogramming of melanomas can be achieved by treatment with both these agents; treatment with either agent alone only achieves reversible differentiation. Genetic variation in this gene is associated with diabetes mellitus insulin-depend

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images