



SEC61B Polyclona Antibody

Catalog No	BYab-10891
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	IHC; ELISA
Gene Name	SEC61B
Protein Name	SEC61B
Immunogen	Synthesized peptide derived from human SEC61B AA range: 30-110
Specificity	This antibody detects endogenous levels of human SEC61B
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:50-200, ELISA(peptide)1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Protein transport protein Sec61 subunit beta
Observed Band	
Cell Pathway	Endoplasmic reticulum membrane ; Single-pass membrane protein .
Tissue Specificity	Brain,Embryonic kidney,Epithelium,Stomach mucosa,
Function	function:Necessary for protein translocation in the endoplasmic reticulum.,similarity:Belongs to the SEC61-beta family.,subunit:Heterotrimeric complex composed of SEC61-alpha, SEC61-beta and SEC61-gamma. Part of a complex composed of SEC61, SEC62 and SEC63. Interacts with SEC62.,
Background	The Sec61 complex is the central component of the protein translocation apparatus of the endoplasmic reticulum (ER) membrane. Oligomers of the Sec61 complex form a transmembrane channel where proteins are translocated across and integrated into the ER membrane. This complex consists of three membrane proteins- alpha, beta, and gamma. This gene encodes the beta-subunit protein. The Sec61 subunits are also observed in the post-ER compartment, suggesting that these proteins can escape the ER and recycle back. There is evidence for

Nanjing BYabscience technology Co.,Ltd



multiple polyadenylated sites for this transcript. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).