



DDA1 rabbit pAb

Catalog No	BYab-11668
Isotype	IgG
Reactivity	Human; Mouse
Applications	IHC;IF
Gene Name	DDA1 C19orf58 PCIA1
Protein Name	DDA1
Immunogen	Synthesized peptide derived from human DDA1 AA range: 49-99
Specificity	This antibody detects endogenous levels of DDA1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	
Tissue Specificity	
Function	function:May be involved in ubiquitination and subsequent proteasomal degradation of target proteins. Component of the DDD-E2 complexes which may provide a platform for interaction with CUL4A and WD repeat proteins.,similarity:Belongs to the DDA1 family.,subunit:Part of the DDD core complex containing DET1, DDA1 and DDB1. The DDD core complex recruits a specific UBE2E enzyme, such as UBE2E1, UBE2E2 UBE2E3, to form specific DDD-E2 complexes.,
Background	

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).