



# RBFA rabbit pAb

|                           |                                                                                                                                   |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| Catalog No                | BYab-11732                                                                                                                        |
| Isotype                   | IgG                                                                                                                               |
| Reactivity                | Human; Mouse                                                                                                                      |
| Applications              | WB;IHC                                                                                                                            |
| Gene Name                 | RBFA C18orf22                                                                                                                     |
| Protein Name              | RBFA                                                                                                                              |
| Immunogen                 | Synthesized peptide derived from human RBFA AA range: 286-336                                                                     |
| Specificity               | This antibody detects endogenous levels of RBFA at Human/Mouse                                                                    |
| Formulation               | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                           |
| Source                    | Polyclonal, Rabbit,IgG                                                                                                            |
| Purification              | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.                         |
| Dilution                  | WB 1:500-2000;IHC-p 1:50-300                                                                                                      |
| Concentration             | 1 mg/ml                                                                                                                           |
| Purity                    | ≥90%                                                                                                                              |
| Storage Stability         | -20°C/1 year                                                                                                                      |
| Synonyms                  |                                                                                                                                   |
| Observed Band             |                                                                                                                                   |
| Cell Pathway              | Mitochondrion .                                                                                                                   |
| Tissue Specificity        |                                                                                                                                   |
| Function                  | similarity:Belongs to the rbfA family.,                                                                                           |
| Background                |                                                                                                                                   |
| matters needing attention | Avoid repeated freezing and thawing!                                                                                              |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Nanjing BYabscience technology Co.,Ltd



## Products Images

Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4° over night

Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).