



# KCTD5 rabbit pAb

|                           |                                                                                                                                                                                                                        |
|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog No                | BYab-11743                                                                                                                                                                                                             |
| Isotype                   | IgG                                                                                                                                                                                                                    |
| Reactivity                | Human; Mouse;Rat                                                                                                                                                                                                       |
| Applications              | WB;ELISA;IHC                                                                                                                                                                                                           |
| Gene Name                 | KCTD5                                                                                                                                                                                                                  |
| Protein Name              | KCTD5                                                                                                                                                                                                                  |
| Immunogen                 | Synthesized peptide derived from human KCTD5 AA range: 170-220                                                                                                                                                         |
| Specificity               | This antibody detects endogenous levels of KCTD5 at Human/Mouse/Rat                                                                                                                                                    |
| Formulation               | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                |
| Source                    | Polyclonal, Rabbit,IgG                                                                                                                                                                                                 |
| Purification              | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.                                                                                                              |
| Dilution                  | WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000                                                                                                                                                                         |
| Concentration             | 1 mg/ml                                                                                                                                                                                                                |
| Purity                    | ≥90%                                                                                                                                                                                                                   |
| Storage Stability         | -20°C/1 year                                                                                                                                                                                                           |
| Synonyms                  |                                                                                                                                                                                                                        |
| Observed Band             |                                                                                                                                                                                                                        |
| Cell Pathway              | Cytoplasm, cytosol . Cytoplasm . Nucleus. Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep proteins. .                                                                                 |
| Tissue Specificity        |                                                                                                                                                                                                                        |
| Function                  | similarity:Contains 1 BTB (POZ) domain.,subcellular location:Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep proteins.,subunit:Interacts with adeno-associated virus 2 REP proteins., |
| Background                |                                                                                                                                                                                                                        |
| matters needing attention | Avoid repeated freezing and thawing!                                                                                                                                                                                   |

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658



#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

#### Products Images

Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night

Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).