



ZN521 rabbit pAb

Catalog No	BYab-11765
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	ZNF521 EHZF LIP3
Protein Name	ZN521
Immunogen	Synthesized peptide derived from human ZN521 AA range: 997-1047
Specificity	This antibody detects endogenous levels of ZN521 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus .
Tissue Specificity	Predominantly expressed in hematopoietic cells. Present in organs and tissues that contain stem and progenitor cells, myeloid and/or lymphoid: placenta, spleen, lymph nodes, thymus, bone marrow and fetal liver. Within the hematopoietic system, it is abundant in CD34(+) cells but undetectable in mature peripheral blood leukocytes, and its levels rapidly decrease during the differentiation of CD34(+) cells in response to hemopoietins.
Function	disease:A chromosomal aberration involving ZNF521 is found in acute lymphoblastic leukemia. Translocation t(9;18)(p13;q11.2) with PAX5. The translocation generates the PAX5-ZNF521 oncogene consisting of the N-terminus part of PAX5 and the C-terminus part of ZNF521.,domain:Uses different DNA- and protein-binding zinc fingers to regulate the distinct BMP-Smad and hematopoietic system.,function:Transcription factor that can both act as an activator or a repressor depending on the context. Involved in BMP signaling and in the regulation of the immature compartment of the hematopoietic system.

Nanjing BYabscience technology Co.,Ltd



Associates with SMADs in response to BMP2 leading to activate transcription of BMP target genes. Acts as a transcriptional repressor via its interaction with EBF1, a transcription factor involved specification of B-cell lineage; this interaction preventing EBF1 to bind DNA and activate target genes.,sim

Background


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night